

AN ANALOGUE OF LEBESGUE-RADON-NIKODYM THEOREM WITH RESPECT TO p -ADIC q -INVARIANT DISTRIBUTION ON $\mathbb{Z}_p$

TAEKYUN KIM

*Institute of Science Education,
Kongju National University, Kongju 314-701, S. Korea
e-mail: tkim64@hanmail.net (or tkim@kongju.ac.kr)*

ABSTRACT. The purpose of this paper is to derive the analogue of Lebesgue-Radon-Nikodym theorem with respect to p -adic q -invariant distribution on $\mathbb{Z}_p$ which is defined by author in [1].

§1. INTRODUCTION

Let p be a fixed prime. Throughout this paper $\mathbb{Z}$, $\mathbb{Z}_p$, $\mathbb{Q}_p$, and $\mathbb{C}_p$ will, respectively, denote the ring of rational integers, the ring of p -adic rational integers, the field of p -adic rational numbers and the completion of algebraic closure of $\mathbb{Q}_p$, cf.[1, 2, 3]. Let v_p be the normalized exponential valuation of $\mathbb{C}_p$ with $|p| = p^{-v_p(p)} = p^{-1}$ and let $a + p^N\mathbb{Z}_p = \{x \in \mathbb{Z}_p | x \equiv a \pmod{p^N}\}$, where $a \in \mathbb{Z}$ lies in $0 \leq a < p^N$. In this paper we assume that $q \in \mathbb{C}_p$ with $|1 - q| < p^{-\frac{1}{p-1}}$ as an indeterminate. We now use the notation

$$[x]_q = [x : q] = \frac{1 - q^x}{1 - q}, \text{ cf. [1, 2, 3].}$$

For any positive integer N we set

$$(1) \quad \mu_q(a + p^N\mathbb{Z}_p) = \frac{q^a}{[p^N]_q}, \text{ (see [1]),}$$

Key words and phrases. p -adic q -invariant measure, Lebesgue decomposition, q -Volkenborn integral, Lebesgue-Radon-Nikodym theorem Bernoulli numbers.

2000 Mathematics Subject Classification: 11S80, 11B68, 11M99 .

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ -TEX

and this can be extended to a distribution on $\mathbb{Z}_p$. In this paper μ_q will be called p -adic q -invariant distribution on $\mathbb{Z}_p$.

For $f \in C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p) = \{f|f : \mathbb{Z}_p \rightarrow \mathbb{C}_p \text{ is } C^{(1)}\text{-function}\}$, the above distribution μ_q yields an integral as follows:

$$(2) \quad I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \quad (\text{ see [1] }) .$$

This integral is called q -Volkenborn integration, cf.[1]. In [1, 3], the q -Bernoulli numbers are defined as

$$(3) \quad \beta_{m,q} = \int_{\mathbb{Z}_p} q^{-x} [x]_q^m d\mu_q(x), \quad m \geq 0 .$$

Note that

$$\beta_{0,q} = \frac{q-1}{\log q}, \quad \beta_{m,q} = \frac{1}{(1-q)^m} \sum_{i=0}^{\infty} \binom{m}{i} (-1)^{m-i} \frac{i}{[i]_q}, \quad \text{cf. [1]}.$$

Also, the q -Bernoulli polynomials are defined by

$$(4) \quad \beta_{n,q}(x) = \int_{\mathbb{Z}_p} q^{-y} [x+y]_q^n d\mu_q(y) = \sum_{k=0}^n \binom{n}{k} q^{kx} \beta_{k,q} [x]_q^{n-k}, \quad n \geq 0.$$

Let $C(\mathbb{Z}_p, \mathbb{C}_p)$ be the space of continuous function on $\mathbb{Z}_p$ with values in $\mathbb{C}_p$, provided with norm $\|f\|_{\infty} = \sup_{x \in \mathbb{Z}_p} |f(x)|$. The difference quotient $\Delta_1 f$ of f is the function of two variables given by

$$\Delta_1 f(m, x) = \frac{f(x+m) - f(x)}{m}, \quad \text{for all } x, m \in \mathbb{Z}_p, m \neq 0.$$

A function $f : \mathbb{Z}_p \rightarrow \mathbb{C}_p$ is said to be a Lipschitz function if there exists a constant $M > 0$ (the Lipschitz constant of f) such that

$$|\Delta_1 f(m, x)| \leq M \text{ for all } m \in \mathbb{Z}_p \setminus \{0\} \text{ and } x \in \mathbb{Z}_p.$$

The $\mathbb{C}_p$ -linear space consisting of all Lipschitz function (or $C^{(1)}$ -function) is denoted by $Lip(\mathbb{Z}_p, \mathbb{C}_p)$ (or $C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$). This space is a Banach space with respect to the norm $\|f\|_1 = \|f\|_{\infty} \vee \|\Delta_1 f\|_{\infty}$. The purpose of this paper is to derive the analogue of Lebesgue-Radon-Nikodym theorem with respect to p -adic q -invariant distribution on $\mathbb{Z}_p$ which is defined by author in [1].

2. ANALOGUE OF LEBESGUE-RADON-NIKODYM THEOREM WITH RESPECT TO μ_q ON $\mathbb{Z}_p$

For $f \in C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$, the q -Volkenborn integral is defined by

$$I_q(f) = \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \frac{1}{[p^N]_q} \sum_{x=0}^{p^N-1} f(x) q^x, \text{ cf. [1].}$$

This definition is based on the p -adic q -distribution on $\mathbb{Z}_p$, defined by $\mu_q(a + p^N \mathbb{Z}_p) = \frac{q^a}{[p^N]_q}$. By the meaning of the extension of q -Volkenborn integral, we consider the below weakly (strongly) p -adic q -invariant distribution μ_q on $\mathbb{Z}_p$, satisfying

$$(5) \quad |[p^n]_q \mu_q(a + p^n \mathbb{Z}_p) - [p^{n+1}]_q \mu_q(a + p^{n+1} \mathbb{Z}_p)| \leq \delta_n,$$

where $\delta_n \rightarrow 0$, $a \in \mathbb{Z}$, and δ_n is independent of a (for strongly p -adic q -distribution, δ_n is replaced by cp^{-n} , where c is positive real constant). Let $f(x)$ be a function on $\mathbb{Z}_p$. Then the q -Volkenborn integral of f with respect to weakly μ_q is

$$(6) \quad \int_{\mathbb{Z}_p} f(x) d\mu_q(x) = \lim_{N \rightarrow \infty} \sum_{x=0}^{p^N-1} f(x) \mu_q(x + p^N \mathbb{Z}_p),$$

if the limit exists. In this case we say that f is q -Volkenborn integrable with respect to μ_q . A distribution μ_q on $\mathbb{Z}_p$ is called 1-admissible if

$$|\mu_q(a + p^N \mathbb{Z}_p)| \leq \frac{1}{|[p^N]_q|} c_n,$$

where $c_n \rightarrow 0$. If μ_q is 1-admissible on $\mathbb{Z}_p$, then we see that μ_q is a weakly p -adic q -invariant distribution on $\mathbb{Z}_p$. Let μ_q be a weakly p -adic q -invariant distribution on $\mathbb{Z}_p$, then $\lim_N [p^N]_q \mu_q(x + p^N \mathbb{Z}_p)$ exists for all $x \in \mathbb{Z}_p$, and converges uniformly with respect to x . We define the q -analogue of Radon-Nikodym derivative of μ_q with respect to p -adic q -invariant distribution on $\mathbb{Z}_p$ as follows:

$$(7) \quad f_{\mu_q}(x) = \lim_{N \rightarrow \infty} [p^N]_q \mu_q(x + p^N \mathbb{Z}_p).$$

Let $x, y \in \mathbb{Z}_p$ with $|x - y|_p = p^{-m}$. Then $x + p^m \mathbb{Z}_p = y + p^m \mathbb{Z}_p$. Hence, we note that

$$(8) \quad |f_{\mu_q}(x) - f_{\mu_q}(y)| = |f_{\mu_q}(x) - [p^n]_q \mu_q(x + p^n \mathbb{Z}_p) - f_{\mu_q}(y) + [p^n]_q \mu_q(y + p^n \mathbb{Z}_p) + [p^n]_q \mu_q(x + p^n \mathbb{Z}_p) - [p^m]_q \mu_q(x + p^m \mathbb{Z}_p) - [p^n]_q \mu_q(y + p^n \mathbb{Z}_p) + [p^m]_q \mu_q(y + p^m \mathbb{Z}_p)|.$$

If μ_q is strongly p -adic q -invariant distribution on $\mathbb{Z}_p$, then we see that

$$(9) \quad |[p^n]_q \mu_q(x + p^n \mathbb{Z}_p) - [p^m]_q \mu_q(x + p^m \mathbb{Z}_p)| \leq Cp^{-m},$$

for some constant C , and $m < n$.

For $n \gg 0$, we note that

$$|f_{\mu_q}(x) - f_{\mu_q}(y)| \leq C_1 p^{-m} = C_1 |x - y|,$$

where C_1 is positive real constant.

Therefore we obtain the following proposition :

Proposition 1. *Let μ_q be a strongly p -adic q -invariant distribution on $\mathbb{Z}_p$. Then $f_{\mu_q} \in Lip(\mathbb{Z}_p, \mathbb{C}_p)$.*

Let $f \in C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$. For any positive integer a , with $a < p^N$, define

$$(10) \quad \mu_{f,q}(a + p^n \mathbb{Z}_p) = \int_{a+p^n \mathbb{Z}_p} q^{-xp^n-a} f(x) d\mu_q(x),$$

where the integral is q -Volkenborn integral.

Note that

$$\begin{aligned} \mu_{f,q}(a + p^n \mathbb{Z}_p) &= \lim_{m \rightarrow \infty} \frac{1}{[p^{m+n}]_q} \sum_{x=0}^{p^m-1} f(a + p^n x) = \lim_{m \rightarrow \infty} \frac{1}{[p^m]_q} \sum_{x=0}^{p^{m-n}-1} f(a + p^n x) \\ &= \frac{1}{[p^n]_q} \int_{\mathbb{Z}_p} q^{-p^n x} f(a + p^n x) d\mu_{q^{p^n}}(x). \end{aligned}$$

By(10), we obtain the following proposition:

Proposition 2. *For $f, g \in C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$, we have*

$$(11) \quad \begin{aligned} \mu_{\alpha f + \beta g, q} &= \alpha \mu_{f, q} + \beta \mu_{g, q}, \\ |\mu_{f, q}(a + p^n \mathbb{Z}_p)| &\leq \|f\|_1 \left| \frac{1}{[p^n]_q} \right|. \end{aligned}$$

Let $P(x) \in \mathbb{C}_p[[x]_q]$ be an arbitrary polynomials. We now give the proof that $\mu_{P,q}$ is a strongly p -adic q -invariant distribution on $\mathbb{Z}_p$. It is enough to prove the statement for $P(x) = [x]_q^k$. Let a be an integer with $0 \leq a < p^n$. Then we see that

$$\begin{aligned} \mu_{P,q}(a + p^n \mathbb{Z}_p) &= \lim_m \frac{1}{[p^m]_q} \sum_{i=0}^{p^{m-n}-1} [a + ip^n]_q^k \\ &= \lim_{m \rightarrow \infty} \frac{1}{[p^m]_q} (p^{m-n} [a]_q^k + k [a]_q^{k-1} q^a [p^n]_q \sum_{i=0}^{p^{m-n}-1} [i]_{q^{p^n}} + \cdots + [p^n]_q^k q^{ak} \sum_{i=0}^{p^{m-n}-1} [i]_{q^{p^n}}^k). \end{aligned}$$

Because $[a + ip^n]_q = ([a]_q + q^a [p^n]_q [i]_{q^{p^n}})$. From (3), we note that

$$\beta_{n,q} = \int_{\mathbb{Z}_p} q^{-x} [x]_q^n d\mu_q(x) = \lim_{m \rightarrow \infty} \frac{1}{[p^m]_q} \sum_{i=0}^{p^m-1} [i]_q^n.$$

Thus, we have

$$\begin{aligned} (12) \quad & [p^n]_q \mu_{P,q}(a + p^n \mathbb{Z}_p) \\ &= -\frac{1 - q^{p^n}}{p^n \log q} [a]_q^k + [p^n]_q q^a \binom{k}{1} [a]_q^{k-1} \beta_{1,q^{p^n}} + \cdots + [p^n]_q^k q^{ak} \beta_{k,q^{p^n}} \\ &= \sum_{i=0}^{\infty} \frac{P^{(i)}(a)}{i!} \beta_{i,q^{p^n}} [p^n]_q^i q^{ai}, \text{ where } P^{(i)}(a) = \left(\frac{d}{d[x]_q} \right)^i P(x)|_{x=a}. \end{aligned}$$

The Eq.(12) can be rewritten as

$$(13) \quad \mu_{P,q}(a + p^n \mathbb{Z}_p) \equiv \frac{P(a)}{[p^n]_q} \beta_{0,q^{p^n}} + q^a \beta_{1,q^{p^n}} P'(a) \pmod{[p^n]_q}.$$

Let x be an arbitrary in $\mathbb{Z}_p$, $x \equiv x_n \pmod{p^n}$, $x \equiv x_{n+1} \pmod{p^{n+1}}$, where x_n, x_{n+1} are positive integers such that $0 \leq x_n < p^n$ and $0 \leq x_{n+1} < p^{n+1}$. Then

$$\begin{aligned} (14) \quad & |[p^n]_q \mu_{P,q}(x + p^n \mathbb{Z}_p) - [p^{n+1}]_q \mu_{P,q}(x + p^{n+1} \mathbb{Z}_p)| \\ &= \left| \sum_{i=0}^{\infty} \frac{P^{(i)}(x_n)}{i!} \beta_{i,q^{p^n}} [p^n]_q^i q^{ai} - \sum_{i=0}^{\infty} \frac{P^{(i)}(x_{n+1})}{i!} \beta_{i,(q^p)^{p^n}} [p^n]_{q^p}^i [p]_q^i q^{ai} \right|. \end{aligned}$$

By (3), (8), (13) and (14), we easily see that

$$|[p^n]_q \mu_{P,q}(x + p^n \mathbb{Z}_p) - [p^{n+1}]_q \mu_{P,q}(x + p^{n+1} \mathbb{Z}_p)| \leq C_2 p^{-n}, \text{ for some constant } C_2.$$

Thus, we note that

$$\begin{aligned} f_{\mu_{P,q}}(a) &= \lim_n [p^n]_q \mu_{P,q}(a + p^n \mathbb{Z}_p) \\ &= \lim_{n \rightarrow \infty} \left(\frac{q^{p^n} - 1}{\log q^{p^n}} [a]_q^k + [p^n]_q q^a [a]_q^{k-1} \binom{k}{1} \beta_{1,q^{p^n}} + \cdots + [p^n]_q^k q^{ka} \beta_{k,q^{p^n}} \right) = [a]_q^k = P(a), \end{aligned}$$

because $\lim_{n \rightarrow \infty} q^{p^n} = 1$ for $|1 - q|_p < p^{-\frac{1}{p-1}}$.

For all $x \in \mathbb{Z}_p$, we have $f_{\mu_{P,q}}(x) = P(x)$, since $f_{\mu_{P,q}}(x)$ is continuous in (7). Now let $g \in C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$. By (12), we easily see that

$$\begin{aligned} &\int_{\mathbb{Z}_p} g(x) d\mu_{P,q}(x) \\ &= \lim_n \sum_{i=0}^{p^n-1} g(i) \mu_{P,q}(i + p^n \mathbb{Z}_p) = \lim_n \frac{1}{[p^n]_q} \sum_{i=0}^{p^n-1} g(i) \sum_{j=0}^k \binom{k}{j} \beta_{j,q^{p^n}} [p^n]_q^j q^{ij} [i]_q^{k-j} \\ &= \sum_{j=0}^k \binom{k}{j} \left(\lim_n \beta_{j,q^{p^n}} \frac{[p^n]_q^j}{[p^n]_q} \sum_{i=0}^{p^n-1} g(i) q^{ij} [i]_q^{k-j} \right) = \lim_n \beta_{0,q^{p^n}} \frac{1}{[p^n]_q} \sum_{i=0}^{p^n-1} g(i) [i]_q^k \\ &= \int_{\mathbb{Z}_p} g(x) [x]_q^k q^{-x} d\mu_q(x), \end{aligned}$$

where the last integral is q -Volkenborn integral on $\mathbb{Z}_p$ which is defined by author in [1]. Therefore we obtain the following theorem:

Theorem 3. *Let $P(x) \in \mathbb{C}_p[[x]_q]$ be an arbitrary polynomials. Then $\mu_{P,q}$ is a strongly p -adic q -invariant distribution, and for all $x \in \mathbb{Z}_p$,*

$$(15) \quad f_{\mu_{P,q}}(x) = P(x).$$

Furthermore, for any $g \in C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$,

$$(16) \quad \int_{\mathbb{Z}_p} g(x) d\mu_{P,q}(x) = \int_{\mathbb{Z}_p} g(x) P(x) q^{-x} d\mu_q(x),$$

where the second integral is q -Volkenborn integral.

Let $f(x) = \sum_{n=0}^{\infty} a_{n,q} \binom{x}{n}_q$ be the q -Mahler expansion of the $C^{(1)}$ -function of f . Here, $\binom{x}{n}_q = \frac{[x]_q[x-1]_q \cdots [x-n+1]_q}{[n]_q[n-1]_q \cdots [1]_q}$, cf.[2]. Then we note that $\lim_n n|a_{n,q}| = 0$, (see [2]). Remark. In the recent, the q -Mahler expansion was introduced by K. Conrad, cf.[2].

Consider $f_m(x) = \sum_{i=0}^m a_{i,q} \binom{x}{i}_q \in \mathbb{C}_p[[x]_q]$. Then $\|f - f_m\|_1 \leq \sup_{n \geq m} n|a_{n,q}|$, cf.[1, 2]. Writing $f = f_m + (f - f_m)$. By using (11), we can observe that

$$\begin{aligned} & |[p^n]_q \mu_{f,q}(a + p^n \mathbb{Z}_p) - [p^{n+1}]_q \mu_{f,q}(a + p^{n+1} \mathbb{Z}_p)| \\ & \leq \max\{|[p^n]_q \mu_{f_m,q}(a + p^n \mathbb{Z}_p) - [p^{n+1}]_q \mu_{f_m,q}(a + p^{n+1} \mathbb{Z}_p)|, \\ & |[p^n]_q \mu_{f-f_m,q}(a + p^n \mathbb{Z}_p) - [p^{n+1}]_q \mu_{f-f_m,q}(a + p^{n+1} \mathbb{Z}_p)|\}. \end{aligned}$$

From Theorem 3, we note that $\mu_{f_m,q}$ is a strongly p -adic q -invariant distribution on $\mathbb{Z}_p$. Moreover, if $m \gg 0$, then we know that

$$|[p^n]_q \mu_{f-f_m,q}(a + p^n \mathbb{Z}_p)| \leq \|f - f_m\|_1 \leq C_3 p^{-n},$$

where C_3 is some positive real constant. Also for $m \gg 0$, it follows that $\|f\|_1 = \|f_m\|_1$ and so

$$|[p^n]_q \mu_{f_m,q}(a + p^n \mathbb{Z}_p) - [p^{n+1}]_q \mu_{f_m,q}(a + p^{n+1} \mathbb{Z}_p)| \leq C_4 p^{-n},$$

where C_4 is also some positive real constant.

We now observe that

$$\begin{aligned} & |f(a) - [p^n]_q \mu_{f,q}(a + p^n \mathbb{Z}_p)| \\ & = |f(a) - f_m(a) + f_m(a) - [p^n]_q \mu_{f_m,q}(a + p^n \mathbb{Z}_p) - [p^n]_q \mu_{f-f_m,q}(a + p^n \mathbb{Z}_p)| \\ & \leq \max\{|f(a) - f_m(a)|, |f_m(a) - [p^n]_q \mu_{f_m,q}(a + p^n \mathbb{Z}_p)|, |[p^n]_q \mu_{f-f_m,q}(a + p^n \mathbb{Z}_p)|\} \\ & \leq \max\{|f(a) - f_m(a)|, |f_m(a) - [p^n]_q \mu_{f_m,q}(a + p^n \mathbb{Z}_p)|, \|f - f_m\|_1\}. \end{aligned}$$

Now if we fix $\epsilon > 0$, and fix m such that $\|f - f_m\|_1 \leq \epsilon$, then for $n \gg 0$, we have

$$|f(a) - [p^n]_q \mu_{f,q}(a + p^n \mathbb{Z}_p)| \leq \epsilon.$$

That is,

$$(17) \quad f_{\mu_{f,q}}(a) = \lim_{n \rightarrow \infty} [p^n]_q \mu_{f,q}(a + p^n \mathbb{Z}_p) = f(a).$$

Let m be the sufficiently large number such that $\|f - f_m\|_1 \leq p^{-2n}$. Then we see that

$$|\mu_{f-f_m, q}(a + p^n \mathbb{Z}_p)| \leq \frac{1}{|[p^n]_q|} \|f - f_m\|_1 \leq [p^n]_q p^{-2n} \leq p^{-n}.$$

Hence, we have

$$(18) \quad \begin{aligned} \mu_{f, q}(a + p^n \mathbb{Z}_p) &= \mu_{f_m, q}(a + p^n \mathbb{Z}_p) + \mu_{f-f_m, q}(a + p^n \mathbb{Z}_p) \equiv \mu_{f_m, q}(a + p^n \mathbb{Z}_p) \\ &\equiv \frac{f_m(a)}{[p^n]_q} \beta_{0, q^{p^n}} - \beta_{1, q^{p^n}} f'_m(a) = \frac{f(a)}{[p^n]_q} \beta_{0, q^{p^n}} - \beta_{1, q^{p^n}} f'(a) \pmod{[p^n]_q}. \end{aligned}$$

Let $g \in C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$. Then

$$\begin{aligned} \int_{\mathbb{Z}_p} g(x) d\mu_{f, q}(x) &= \lim_n \sum_{i=0}^{p^n-1} g(i) \mu_{f, q}(i + p^n \mathbb{Z}_p) \\ &= \lim_n \sum_{i=0}^{p^n-1} g(i) \left(\frac{f(i)}{[p^n]_q} \beta_{0, q^{p^n}} - \beta_{1, q^{p^n}} f'(i) \right) \\ &= \lim_n \left(\frac{1}{[p^n]_q} \sum_{i=0}^{p^n-1} g(i) f(i) - \frac{1}{2} \sum_{i=0}^{p^n-1} g(i) f'(i) \right) \\ &= \int_{\mathbb{Z}_p} g(x) f(x) q^{-x} d\mu_q(x), \end{aligned}$$

because $\lim_{n \rightarrow \infty} \beta_{k, q^n} = B_k$, where B_k are the ordinary k -th Bernoulli numbers.

Let δ be the function from $C^{(1)}(\mathbb{Z}_p, \mathbb{C}_p)$ to $Lip(\mathbb{Z}_p, \mathbb{C}_p)$. We know that μ_q is a strongly p -adic q -invariant distribution on $\mathbb{Z}_p$, then $|f_{\mu_q}(a) - [p^n]_q \mu_q(a + p^n \mathbb{Z}_p)| \leq C p^{-n}$ for any positive integer n . Let $\mu_{1, q}$ be the strongly p -adic q -invariant distribution associated to f_{μ_q} ($\mu_{1, q} = \delta(f_{\mu_q})$). If $\mu_{1, q}$ is the associated strongly p -adic q -invariant distribution on $\mathbb{Z}_p$, then, by (18), we have

$$|[p^n]_q \mu_{1, q}(a + p^n \mathbb{Z}_p) - f_{\mu_q}(a)| \leq \max\{p^{-n}, p^{-n} |\beta_{0, q^{p^n}} f'(a)|\} \leq C_5 p^{-n}, \text{ for } n \gg 0,$$

where C_5 is a some positive real constant. Because $[p^n]_q \equiv 0 \pmod{p^n}$.

This shows that for sufficient large n

$$\begin{aligned} &|\mu_q(a + p^n \mathbb{Z}_p) - \mu_{1, q}(a + p^n \mathbb{Z}_p)| \\ &= \frac{1}{|[p^n]_q|} |[p^n]_q \mu_q(a + p^n \mathbb{Z}_p) - f_{\mu_q}(a) + f_{\mu_q}(a) - [p^n]_q \mu_{1, q}(a + p^n \mathbb{Z}_p)| \\ &\leq \max\{p^n p^{-n} C_6, p^n p^{-n} C_7\} \leq M, \end{aligned}$$

where C_6, C_7, M are some positive real constants.

Thus, $\mu_q - \mu_{1,q}$ is a distribution and bounded. Hence, $\mu_q - \mu_{1,q}$ is a measure on $\mathbb{Z}_p$. Therefore we obtain the following theorem:

Theorem 4. *Let μ_q be a strongly p -adic q -invariant distribution on $\mathbb{Z}_p$, and assume that the Radon-Nikodym derivative f_{μ_q} on $\mathbb{Z}_p$ is a $C^{(1)}$ -function. Suppose that $\mu_{1,q}$ is the strongly p -adic q -invariant distribution associated to f_{μ_q} ($\mu_{1,q} = \delta(f_{\mu_q})$), then there exists a measure $\mu_{2,q}$ on $\mathbb{Z}_p$ such that*

$$(19) \quad \mu_q = \mu_{1,q} + \mu_{2,q}.$$

Eq.(19) is p -adic analogue of Lebesgue decomposition with respect to strongly p -adic q -invariant distribution on $\mathbb{Z}_p$.

REFERENCES

1. T. Kim, *q-Volkenborn integration*, Russ. J. Math. Phys. **9** (2002), 288-299.
2. T. Kim, S. D. Kim, D. W. Park, *On uniform differentiability and q -Mahler expansion*, Advan. Stud. Contemp. Math. **4 no. 1** (2001), 35-41.
3. T. Kim, *An invariant p -adic integral associated with Daehee numbers*, Integral Trans. Special Funct. **13** (2002), 65-69.